

# Charged, rotating black objects in Einstein-Maxwell-dilaton theory in $D \geq 5$

Burkhard Kleihaus, Jutta Kunz<sup>†</sup> and Eugen Radu<sup>‡</sup>

<sup>†</sup>Institut für Physik, Universität Oldenburg, Postfach 2503 D-26111 Oldenburg, Germany

<sup>‡</sup>Departamento de Física da Universidade de Aveiro and CIDMA,  
Campus de Santiago, 3810-183 Aveiro, Portugal

May 20, 2016

## Abstract

We show that the general framework proposed in [1] for the study of asymptotically flat vacuum black objects with  $k + 1$  equal magnitude angular momenta in  $D \geq 5$  spacetime dimensions (with  $0 \leq k \leq [\frac{D-5}{2}]$ ) can be extended to the case of Einstein-Maxwell-dilaton (EMd) theory. This framework can describe black holes with spherical horizon topology, the simplest solutions corresponding to a class of electrically charged (dilatonic) Myers-Perry black holes. Balanced charged black objects with  $S^{n+1} \times S^{2k+1}$  horizon topology can also be studied (with  $D = 2k + n + 4$ ). Black rings correspond to the case  $k = 0$ , while the solutions with  $k > 0$  are *black ringoids*. The basic properties of EMd solutions are discussed for the special case of a Kaluza-Klein value of the dilaton coupling constant. We argue that all features of these solutions can be derived from those of the vacuum seed configurations.

## 1 Introduction

The study of black hole (BH) solutions in more than  $D = 4$  dimensions is a subject of long standing interest in General Relativity<sup>1</sup>. A seminal result in this area was the discovery of the  $D = 5$  black ring (BR) by Emparan and Reall [2], [3]. The  $D > 5$  generalizations of the BR were constructed in [4] within an approximation scheme, and fully non-perturbatively in [5], [6] (although for  $D = 6, 7$  only). In contrast to the Myers-Perry (MP) BHs [7], which have a spherical horizon topology being natural higher dimensional generalizations of the  $D = 4$  Kerr solution [8], the BRs have an event horizon of  $S^{D-3} \times S^1$  topology, and possess no four dimensional counterpart.

---

<sup>1</sup>In this work we shall restrict to configurations approaching asymptotically a Minkowski spacetime background.

The rapid developments following the discovery in [2], [3] have revealed the existence of a ‘zoo’ of higher dimensional solutions with various topologies of the event horizon (a review of the existing results can be found in [9], [10], [11]). However, most of the activity in this area concerns the pure Einstein gravity case without matter fields. In particular, to our knowledge, there is no non-perturbative construction of non-vacuum, singularity-free  $D > 5$  black objects<sup>2</sup> with a non-spherical horizon topology<sup>3</sup>.

The main purpose of this work is to propose a general framework for the study of a class of asymptotically flat black objects in Einstein-Maxwell-dilaton (EMd) theory for a number  $D \geq 5$  of spacetime dimensions. These black objects possess  $k + 1$  equal magnitude angular momenta and can describe MP-like BHs with spherical horizon topology or balanced black objects with  $S^{n+1} \times S^{2k+1}$  horizon topology (with  $D = 2k + n + 4$  and  $0 \leq k \leq [\frac{D-5}{2}]$ ). In the absence of matter fields, this framework reduces to that employed in [1] to study BRs ( $k = 0$ ) and black ringoids ( $k > 0$ ). Here we show that the approach in [1] can be extended to the EMd case.

Moreover, for a special value of the dilaton coupling constant, all solutions in [1] can be extended to the EMd case in a straightforward way, by using a generation technique. This approach has the advantage to easily provide a window into the elusive general EMd case; also, we expect some of the solutions’ properties to be generic.

## 2 The framework

### 2.1 The action and field equations

The action of the  $D$ -dimensional EMd theory is ( $G = 1$ )

$$S = \frac{1}{16\pi} \int d^D x \sqrt{-g} \left( R - \frac{1}{2} \Phi_{,\mu} \Phi^{,\mu} - \frac{1}{4} e^{-2a\Phi} F_{\mu\nu} F^{\mu\nu} \right), \quad (1)$$

where  $a$  is the dilaton coupling constant and  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ . The field equations consist of the Einstein equations

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = \frac{1}{2} T_{\mu\nu}, \quad (2)$$

with the stress-energy tensor

$$T_{\mu\nu} = \partial_\mu \Phi \partial_\nu \Phi - \frac{1}{2} g_{\mu\nu} \partial_\tau \Phi \partial^\tau \Phi + e^{-2a\Phi} \left( F_{\mu\tau} F_\nu{}^\tau - \frac{1}{4} g_{\mu\nu} F_{\tau\beta} F^{\tau\beta} \right),$$

the Maxwell equations

$$\nabla_\mu (e^{-2a\Phi} F^{\mu\nu}) = 0, \quad (3)$$

and the dilaton equation

$$\nabla^2 \Phi = -\frac{a}{2} e^{-2a\Phi} F_{\mu\nu} F^{\mu\nu}. \quad (4)$$

---

<sup>2</sup>The situation is different in five dimensions, where a variety of BR solutions with (Abelian) gauge fields and scalars are known in closed form [12] (see also the Einstein-Maxwell numerical solutions in [13]).

<sup>3</sup>Balanced Einstein-Maxwell BHs with  $S^2 \times S^{D-4}$  event horizon topology were constructed in [14]. However, those solutions are not asymptotically flat.

## 2.2 The Ansatz

Following [1], we consider the metric Ansatz

$$ds^2 = f_1(r, \theta) (dr^2 + \Delta(r)d\theta^2) + f_2(r, \theta)d\Omega_n^2 - f_0(r, \theta)dt^2 \\ + f_3(r, \theta)(d\psi + \mathcal{A} - W(r, \theta)dt)^2 + f_4(r, \theta)d\Sigma_k^2, \quad (5)$$

which describes the geometry of black objects with  $k+1$  equal magnitude angular momenta in  $D \geq 5$  spacetime dimensions (with  $D = 2k + n + 4$ ). The above choice of the Ansatz becomes transparent when considering the Minkowski spacetime limit of (5). This background metric is recovered for  $f_0 = f_1 = 1$ ,  $f_2 = r^2 \cos^2 \theta$ ,  $f_3 = f_4 = r^2 \sin^2 \theta$ ,  $W = 0$  and  $\Delta(r) = r^2$ :

$$ds^2 = dr^2 + r^2(d\theta^2 + \cos^2 \theta d\Omega_n^2 + \sin^2 \theta d\Omega_{2k+1}^2) - dt^2, \quad (6)$$

where  $0 \leq r < \infty$ ,  $0 \leq \theta \leq \pi/2$  and  $t$  is the time coordinate. Also,  $d\Omega_n^2$  the metric on the round  $n$ -dimensional sphere, while the metric of a  $(2k+1)$ -dimensional sphere is written as an  $S^1$  fibration over the complex projective space  $\mathbb{CP}^k$ ,

$$d\Omega_{2k+1}^2 = (d\psi + \mathcal{A})^2 + d\Sigma_k^2, \quad (7)$$

where  $d\Sigma_k^2$  is the metric on the unit  $\mathbb{CP}^k$  space and  $\mathcal{A} = \mathcal{A}_i dx^i$  is its Kähler form<sup>4</sup>.

A gauge field Ansatz compatible with the symmetries of the line element (5) reads

$$A = A_t(r, \theta)dt + A_\psi(r, \theta)(d\psi + \mathcal{A}), \quad (8)$$

while the dilaton  $\Phi$  is

$$\Phi = \Phi(r, \theta). \quad (9)$$

## 2.3 Boundary conditions and quantities of interest

In this approach, the dependence of the coordinates on the  $S^{2k+1}$  and  $S^n$  parts of the metric factorizes, such that the problem is effectively codimension-2. As a result, the information on the solutions is encoded in the metric functions  $(f_i, W)$  (with  $i = 0, \dots, 4$ ), the gauge potentials  $(A_t, A_\psi)$  and the dilaton  $\Phi$ . (Note that the function  $\Delta(r)$  which enters (5) is an input ‘background’ function which is chosen for convenience by using the residual metric gauge freedom. The numerical solutions have  $\Delta(r) = r^2$ .)

Then the resulting EMD equations of motion form a set of nine coupled nonlinear partial differential equations (PDEs) in terms of  $(r, \theta)$  only, which are solved subject to the boundary conditions given below<sup>5</sup>.

---

<sup>4</sup>The fibre is parameterized by the coordinate  $\psi$ , which has period  $2\pi$ . Also, the term  $d\Sigma_k^2$  is absent in (5) for  $k = 0$  (in which case  $\mathcal{A} = 0$ ). However, the general relations exhibited below are still valid in that case, see [1].

<sup>5</sup>Note that the metric functions should satisfy a number of extra boundary conditions which guarantee the regularity of the solutions (*e.g.* the constancy of the Hawking temperature on the horizon, see the discussion in [1]).

The range of the  $\theta$ -coordinate is  $[0, \pi/2]$ , while  $r_H \leq r < \infty$ . The event horizon is located at  $r = r_H > 0$ , the metric of a spatial cross-section of the horizon being

$$d\sigma^2 = f_1(r_H, \theta)r_H^2 d\theta^2 + f_2(r_H, \theta)d\Omega_n^2 + f_3(r_H, \theta)(d\psi + \mathcal{A})^2 + f_4(r_H, \theta)d\Sigma_k^2. \quad (10)$$

At  $r = r_H$ , the following boundary conditions are satisfied<sup>6</sup>

$$f_0 = 0, \quad r_H \partial_r f_1 + 2f_1 = \partial_r f_2 = \partial_r f_3 = 0, \quad W = \Omega_H, \quad \partial_r A_\psi = 0, \quad A_t + \Omega_H A_\psi = \Phi_H, \quad \partial_r \Phi = 0. \quad (11)$$

As  $r \rightarrow \infty$ , the Minkowski spacetime background is recovered, with vanishing matter fields, which implies

$$f_0 = f_1 = 1, \quad f_2 = r^2 \cos^2 \theta, \quad f_3 = f_4 = r^2 \sin^2 \theta, \quad W = 0, \quad A_t = A_\psi = \Phi = 0. \quad (12)$$

At  $\theta = \pi/2$ , we impose

$$\partial_\theta f_0 = \partial_\theta f_1 = f_2 = \partial_\theta f_3 = \partial_\theta f_4 = \partial_\theta W = 0, \quad \partial_\theta A_t = A_\psi = 0, \quad \partial_\theta \Phi = 0. \quad (13)$$

The boundary conditions at  $\theta = 0$  are more complicated, depending on the event horizon topology. In the simplest case of solutions with a spherical horizon topology, one imposes

$$\partial_\theta f_0 = \partial_\theta f_1 = \partial_\theta f_2 = f_3 = f_4 = \partial_\theta W = 0, \quad \partial_\theta A_t = A_\psi = 0, \quad \partial_\theta \Phi = 0. \quad (14)$$

However, as discussed at length in [1], the metric Ansatz (5) allows as well for an  $S^{n+1} \times S^{2k+1}$  horizon topology. Such solutions possess a new input parameter  $R_0 > r_H$  (which provides a rough measure for the size of the  $S^{n+1}$  sphere on the horizon), with

$$\partial_\theta f_0 = \partial_\theta f_1 = f_2 = \partial_\theta f_3 = \partial_\theta f_4 = \partial_\theta W = 0, \quad \partial_\theta A_t = \partial_\theta A_\psi = 0, \quad \partial_\theta \Phi = 0, \quad (15)$$

for  $r_H < r \leq R_0$ , while for  $r_H > R_0$  the boundary conditions are given by (14). Thus, for such solutions, the functions  $f_3, f_4$  multiplying the  $S^{2k+1}$  part of the horizon metric (10) are strictly positive and finite for any  $r \leq R_0$ , while  $f_2 \sim \sin^2 2\theta$  and thus vanishes at both  $\theta = 0$  and  $\theta = \pi/2$ . However, the  $S^{n+1}$  and  $S^{2k+1}$  parts in (10) are not round spheres. To obtain a measure for the deformation of the  $S^{n+1}$  sphere, we consider the ratio  $L_e/L_p$ , where  $L_e$  is the circumference at the equator ( $\theta = \pi/4$ , where the sphere is fattest), and  $L_p$  the circumference along the poles,

$$L_e = 2\pi \sqrt{f_2(r_H, \pi/4)}, \quad L_p = 2 \int_0^{\pi/2} d\theta \, r_H \sqrt{f_1(r_H, \theta)}. \quad (16)$$

A possible estimate for the deformation of the sphere  $S^{2k+1}$  in (10) is given by the ratio  $R_{2k+1}^{(in)}/R_{2k+1}^{(out)}$ , where

$$R_{2k+1}^{(in)} = (f_3(r_H, 0)f_4^{2k}(r_H, 0))^{\frac{1}{2(2k+1)}}, \quad R_{2k+1}^{(out)} = (f_3(r_H, \pi/2)f_4^{2k}(r_H, \pi/2))^{\frac{1}{2(2k+1)}}. \quad (17)$$

---

<sup>6</sup>Note that the boundary conditions (11)-(15) are compatible with an approximate form of the solutions on the boundaries of the domain of integration.

The expressions of the event horizon area  $A_H$ , Hawking temperature  $T_H$ , event horizon velocity  $\Omega_H$  and the horizon electrostatic potential  $\Phi_H$  of the solutions are similar for any horizon topology and read

$$A_H = r_H V_{(n)} V_{(2k+1)} \int_0^{\pi/2} d\theta \sqrt{f_1 f_2^n f_3 f_4^{2k}} \Big|_{r=r_H}, \quad (18)$$

$$T_H = \frac{1}{2\pi} \lim_{r \rightarrow r_H} \frac{1}{(r - r_H)} \sqrt{\frac{f_0}{f_1}}, \quad \Omega_H = W|_{r=r_H}, \quad \Phi_H = (A_t + \Omega_H A_\psi)|_{r=r_H},$$

where  $V_{(p)}$  is the area of the unit  $S^p$  sphere. Also, one can see that the Killing vector  $\xi = \partial/\partial_t + \Omega_H \partial/\partial_\psi$  is orthogonal and null on the horizon.

The global charges of the system are the mass  $\mathcal{M}$ , the angular momenta  $J_i$  and the electric charge  $Q_E$ . They are read from the large- $r$  asymptotics of the metric functions and electric potential,  $g_{tt} = -1 + \frac{C_t}{r^{D-3}} + \dots$ ,  $g_{\psi t} = -f_3 W = \frac{C_\psi}{r^{D-3}} \sin^2 \theta + \dots$ ,  $A_t = \frac{Q}{r^{D-3}} + \dots$ , with

$$\mathcal{M} = \frac{(D-2)V_{(D-2)}}{16\pi} C_t, \quad J_1 = \dots = J_{k+1} = \frac{V_{(D-2)}}{8\pi} C_\psi = J, \quad Q_E = \frac{(D-2)V_{(D-2)}}{16\pi} Q. \quad (19)$$

For any horizon topology, these black objects satisfy the Smarr relation

$$(D-3)\mathcal{M} = (D-2)\left(T_H \frac{A_H}{4} + (k+1)\Omega_H J\right) + (D-3)Q_E \Phi_H, \quad (20)$$

and the 1<sup>st</sup> law

$$d\mathcal{M} = \frac{1}{4} T_H dA_H + (k+1)\Omega_H dJ + \Phi_H dQ_E. \quad (21)$$

In the canonical ensemble, we study solutions holding fixed the temperature  $T_H$ , the electric charge  $Q_E$  and the angular momentum  $J$ . The associated thermodynamic potential is the Helmholtz free energy  $F = \mathcal{M} - \frac{1}{4} T_H A_H$ . Black objects in a grand canonical ensemble are also of interest, in which case we keep the temperature  $T_H$ , the chemical potential  $\Phi_H$  and the event horizon velocity  $\Omega_H$  fixed. In this case the thermodynamics is obtained from the Gibbs potential  $G = \mathcal{M} - \frac{1}{4} T_H A_H - (k+1)\Omega_H J - \Phi_H Q_E$ .

Following the usual convention in the literature, we fix the overall scale of the solutions by fixing their mass  $\mathcal{M}$ . Then the solutions are characterized by a set of reduced dimensionless quantities, obtained by dividing out an appropriate power of  $\mathcal{M}$ :

$$j = c_j \frac{J}{\mathcal{M}^{\frac{D-2}{D-3}}}, \quad a_H = c_a \frac{A_H}{\mathcal{M}^{\frac{D-2}{D-3}}}, \quad w_H = c_w \Omega_H \mathcal{M}^{\frac{1}{D-3}}, \quad t_H = c_t T_H \mathcal{M}^{\frac{1}{D-3}}, \quad q = \frac{Q_E}{\mathcal{M}}, \quad (22)$$

with the coefficients

$$\begin{aligned}
c_j &= \frac{(D-2)^{\frac{D-2}{D-3}}}{(16\pi)^{\frac{1}{D-3}} 2^{\frac{D-2}{D-3}}} \frac{1+k}{\sqrt{(D-3)(2k+1)}} (V_{(n+1)} V_{(2k+1)})^{\frac{1}{D-3}}, \\
c_a &= \frac{2^{\frac{2}{D-3}}}{(16\pi)^{\frac{D-2}{D-3}}} (D-2)^{\frac{D-2}{D-3}} \sqrt{\frac{D-2k-4}{D-3}} (V_{(n+1)} V_{(2k+1)})^{\frac{1}{D-3}}, \\
c_w &= \frac{2^{\frac{1}{D-3}}}{(D-2)^{\frac{1}{D-3}}} \sqrt{\frac{D-3}{2k+1}} \frac{(16\pi)^{\frac{1}{D-3}}}{(V_{(n+1)} V_{(2k+1)})^{\frac{1}{D-3}}}, \\
c_t &= \frac{(D-4)\sqrt{D-3}}{2^{\frac{2(D-2)}{D-3}} (D-2)^{\frac{1}{D-3}} (D-2k-4)^{\frac{3}{2}}} \frac{(16\pi)^{\frac{D-2}{D-3}}}{(V_{(n+1)} V_{(2k+1)})^{\frac{1}{D-3}}}.
\end{aligned}$$

For completeness, let us mention that the charged solutions possess also a magnetic moment  $\mu$  and a dilaton charge  $Q_d$ . These quantities are read again from the far field behaviour of the fields,  $A_\psi = -\frac{\mu}{(D-3)V_{(D-2)}r^{D-3}} + \dots$ ,  $\Phi = -\frac{Q_d}{(D-3)V_{(D-2)}r^{D-3}} + \dots$ , and do not enter their thermodynamic description. Also, as usual with charged spinning solutions, a gyromagnetic ratio is defined as

$$g = \frac{2\mu\mathcal{M}}{Q_E J}. \quad (23)$$

### 3 Solutions. The Kaluza-Klein case

The only vacuum solutions which can be written within the Ansatz (5) and are known in closed form are the MP BHs and the  $D = 5$  BR spinning in a single plane. Apart from that, the Refs. [1], [5], [6], [13] gave numerical evidence for the existence of BRs and black ringoids for several values of  $D > 5$ .

Given the above formulation of the problem, EMD generalizations of these configurations can be constructed numerically, by employing the numerical scheme developed in [1] for the vacuum case (see also [15], [16]). Indeed, charged MP BHs were considered in [17], while BR solutions have been studied in Ref. [13], in both cases for  $D = 5$  spacetime dimensions and a pure Einstein-Maxwell theory. By using a similar approach, we have found (preliminary) numerical evidence for the existence of  $D = 7$ ,  $k = 1$  balanced black ringoids, again in the Einstein-Maxwell theory.

However, a numerical investigation of the generic EMD solutions is a complicated task beyond the purposes of this work. In what follows, we shall restrict ourselves to the special case of an EMD model with a Kaluza-Klein value of the dilaton coupling constant  $a$ ,

$$a = \frac{D-1}{\sqrt{2(D-1)(D-2)}}. \quad (24)$$

In this limit, the EMD solutions can be generated by using the (vacuum) Einstein black objects

as seeds<sup>7</sup>. The procedure is well known in the literature and works as follows: we first embed the  $D$ -dimensional vacuum solutions into a  $(D+1)$  spacetime with a trivial extra coordinate  $U$ ,

$$ds_{D+1}^2 = dU^2 + ds^2. \quad (25)$$

Then we perform a boost in the  $t-U$  plane with  $t \rightarrow t \cosh \alpha + U \sinh \alpha$ ,  $U \rightarrow U \cosh \alpha + t \sinh \alpha$ . In the next step we consider the following parametrization of the resulting  $(D+1)$ -dimensional boosted metric

$$ds_{D+1}^2 = e^{\frac{2}{\sqrt{2(D-1)(D-2)}}\Phi} g_{\mu\nu} dx^\mu dx^\nu + e^{-\frac{2(D-2)}{\sqrt{2(D-1)(D-2)}}\Phi} (dU + A_\nu dx^\nu)^2, \quad (26)$$

which allows for a straightforward reduction to  $D$ -dimensions with respect to the Killing vector  $\partial/\partial U$ . Then  $g_{\mu\nu}$ ,  $A_\rho$ , and  $\Phi$  are identified with the  $D$ -dimensional metric, the  $D$ -dimensional Maxwell potential, and the dilaton function, respectively. Also, they satisfy the EMD equations (2)-(4) in  $D$  spacetime dimensions.

Considering a vacuum Einstein gravity solution described by the metric Ansatz (5), a direct computation leads to the following expression of the EMD solution:

$$\begin{aligned} f_0 &= \frac{\left[1 + (1 - f_0^{(0)} + f_3^{(0)} W^{(0)2}) \sinh^2 \alpha\right]^{\frac{1}{D-2}}}{1 + (1 - f_0^{(0)}) \sinh^2 \alpha} f_0^{(0)}, \\ (f_1; f_2; f_4) &= \left[1 + (1 - f_0^{(0)} + f_3^{(0)} W^{(0)2}) \sinh^2 \alpha\right]^{\frac{1}{D-2}} (f_1^{(0)}; f_2^{(0)}; f_4^{(0)}), \\ f_3 &= \frac{1 + (1 - f_0^{(0)}) \sinh^2 \alpha}{\left[1 + (1 - f_0^{(0)} + f_3^{(0)} W^{(0)2}) \sinh^2 \alpha\right]^{\frac{D-3}{D-2}}} f_3^{(0)}, \quad W = \frac{\cosh \alpha}{1 + (1 - f_0^{(0)}) \sinh^2 \alpha} W^{(0)}, \end{aligned} \quad (27)$$

together with

$$\begin{aligned} A_t &= \frac{(1 - f_0^{(0)} + f_3^{(0)} W^{(0)2}) \sinh \alpha \cosh \alpha}{1 + (1 - f_0^{(0)} + f_3^{(0)} W^{(0)2}) \sinh^2 \alpha}, \\ A_\psi &= -\frac{f_3^{(0)} W^{(0)} \sinh \alpha}{1 + (1 - f_0^{(0)} + f_3^{(0)} W^{(0)2}) \sinh^2 \alpha}, \\ \Phi &= -\frac{1}{2(D-2)} \sqrt{2(D-1)(D-2)} \log \left(1 + (1 - f_0^{(0)} + f_3^{(0)} W^{(0)2}) \sinh^2 \alpha\right), \end{aligned} \quad (28)$$

where the superscript (0) stands for the pure Einstein gravity seed metric. One can easily see that these functions satisfy the boundary conditions (11)-(15), since the seed solution is also subject to the same set of conditions.

---

<sup>7</sup>A similar approach has been used in [18], [19] to study the MP BHs in EMD theory with a dilaton coupling constant given by (24).  $D = 5$  BRs and black Saturns have been constructed in [20, 21], again in the same model.




Figure 1: The ratios  $L_e/L_p$  and  $R_{2k+1}^{(in)}/R_{2k+1}^{(out)}$ , which encode the deformation of the horizon, are shown *vs.* the reduced angular momentum  $j$  for  $D = 5, 6$  black ring solutions and  $D = 7$  charged black ringoids in EMD theory. The red curves corresponds to the vacuum solutions. The other curves are for charged solutions with the boosting parameters (from right to left)  $\alpha = 0.5, 1, 1.5$  and  $2$ .

For both the MP BHs and  $D = 5$  BR seed solutions, it is straightforward to write down the corresponding closed form EMD generalizations. For example, in the MP case one replaces in (27)-(28) the following expression of the vacuum seed configuration [1]:

$$\begin{aligned}
 f_0^{(0)} &= \frac{\Delta(r)}{(r^2 + a^2)P(r, \theta)}, & f_1^{(0)} &= \frac{r^2 + a^2 \cos^2 \theta}{\Delta(r)}, & f_2^{(0)} &= r^2 \cos^2 \theta, \\
 f_3^{(0)} &= (r^2 + a^2) \sin^2 \theta P(r, \theta), & f_4^{(0)} &= (r^2 + a^2) \sin^2 \theta, \\
 W^{(0)} &= \frac{M}{r^{D-(2k+5)}} \frac{a}{(r^2 + a^2)^{k+1} (r^2 + a^2 \cos^2 \theta) P(r, \theta)},
 \end{aligned} \tag{29}$$

$M, a$  being two input parameters and

$$\Delta(r) = (r^2 + a^2) \left( 1 - \frac{M}{r^{D-(2k+5)} (r^2 + a^2)^{k+1}} \right), \quad P(r, \theta) = 1 + \frac{M}{r^{D-(2k+5)}} \frac{a^2 \sin^2 \theta}{(r^2 + a^2)^{k+1} (r^2 + a^2 \cos^2 \theta)},$$




Figure 2: The reduced area  $a_H$ , the reduced temperature  $t_H$  and the reduced angular velocity  $w_H$  are shown *vs.* the reduced angular momentum  $j$  for  $D = 5$  charged black rings (BR) and Myers-Perry (MP) black holes.

where a different choice for  $\Delta$  is employed.

Having derived the expressions of the geometry and matter functions, it is straightforward to study all properties of the solutions. For example, in Figure 1 we show the quantities  $L_e/L_p$  and  $R_{2k+1}^{(in)}/R_{2k+1}^{(out)}$  which encode the deformation of the horizon (see (16), (17)) for  $D = 5, 6, 7$  black ring(oid)s and several values of the boosting parameter  $\alpha$ . One can see that the charged solutions share the pattern of the neutral ones, being shifted to smaller values of  $j$ . For example, in the  $D = 5$  case, the hole inside the ring shrinks to zero size while the outer radius goes to infinity as a critical configuration is approached<sup>8</sup>.

<sup>8</sup> All results for MP BHs and  $D = 5$  BRs shown in the plots in this work are found by using the closed form expression of the vacuum seed solutions. For  $D = 5$  BRs, a comparison between the exact solution and the numerically generated one can be found in Appendix B of Ref. [16].

Moreover, for both closed form and numerical solutions, the quantities which enter the first law result from those of the corresponding vacuum seed configurations. A direct computation leads to




Figure 3: Same as Figure 2 for  $D = 6$  dimensions.

$$\mathcal{M} = \left(1 + \frac{D-3}{D-2} \sinh^2 \alpha\right) \mathcal{M}^{(0)}, \quad J = \cosh \alpha \, J^{(0)}, \quad \Omega_H = \frac{1}{\cosh \alpha} \Omega_H^{(0)}, \quad (30)$$

$$T_H = \frac{1}{\cosh \alpha} T_H^{(0)}, \quad A_H = \cosh \alpha \, A_H^{(0)}, \quad Q_E = \frac{D-3}{D-2} \sinh \alpha \cosh \alpha \, \mathcal{M}^{(0)}, \quad \Phi_H = \tanh \alpha,$$

and

$$\begin{aligned}
j &= \left( \frac{D-2}{D-2+(D-3)\sinh^2\alpha} \right)^{\frac{D-2}{D-3}} \cosh\alpha j^{(0)}, \quad a_H = \left( \frac{D-2}{D-2+(D-3)\sinh^2\alpha} \right)^{\frac{D-2}{D-3}} \cosh\alpha a_H^{(0)}, \\
t_H &= \left( 1 + \frac{(D-3)}{(D-2)} \sinh^2\alpha \right)^{\frac{1}{D-3}} \frac{1}{\cosh\alpha} t_H^{(0)}, \quad w_H = \left( 1 + \frac{(D-3)}{(D-2)} \sinh^2\alpha \right)^{\frac{1}{D-3}} \frac{1}{\cosh\alpha} w_H^{(0)}, \\
q &= \frac{(D-3)\sinh\alpha\cosh\alpha}{D-2+(D-3)\sinh^2\alpha},
\end{aligned} \tag{31}$$

for the scaled variables.




Figure 4: Same as Figure 2 for charged black ringoids (Br) and Myers-Perry (MP) black holes in  $D = 7$  dimensions (both with 2 equal magnitude angular momenta).

One can see that the boosting parameter  $\alpha$  is a monotonic function of the horizon electrostatic potential  $\Phi_H$  (or, equally, is uniquely fixed by the reduced charge  $q$ ). Also, given a mass  $\mathcal{M}$ , the electric charge  $Q_E$  cannot be arbitrarily large, with  $q \leq 1$ . The limit  $\alpha \rightarrow \infty$

corresponds to singular black objects, with  $j \rightarrow 0$ ,  $a_H \rightarrow 0$  and  $q \rightarrow 1$ . Moreover, one can show that the Gibbs potential of the charged solutions equals that of the seed vacuum configurations  $G = G^{(0)}$ , while  $F = F^{(0)} + \frac{D-3}{D-2} \sinh^2 \alpha \mathcal{M}^{(0)}$ .

It follows that, for any finite  $\alpha$ , some basic thermodynamic properties of these EMd solutions are qualitatively similar to the vacuum seed case. For example, as shown in Figures 2-4, the  $(j, a_H)$  and  $(j, t_H)$  diagrams of the charged solutions have the same shape for any value of  $q$ . However, the curves in the phase diagram get shifted to lower  $a_H$  and  $j$  as the charge parameter  $q$  is increased.

Also, a generic property of the solutions is the occurrence of a cusp in the  $a_H(j)$  black ring(oid) diagram, where a branch of “fat” black ring(oid) solutions emerges, with the existence of a minimally spinning solution. A comparison of the results (with the set of MP-like solutions included), suggest that, similar to the vacuum case, the  $k \geq 1$  black ringoids with  $S^2 \times S^{2k+1}$  horizon topology are the natural counterparts of the  $D = 5$  BRs. As noticed in [21], the branch of “fat”  $D = 5$  charged BRs ends in a limiting singular solution with  $a_H = 0$  and nonzero  $j$ . The same configuration is also approached by the charged MP BHs with maximal<sup>9</sup>  $j$ . The existing data strongly suggest that this is the picture also for the  $D = 7$  charged Br and MP solutions.




Figure 5: The (area-temperature-charge) diagram is shown for charged Myers-Perry black holes (left) and black rings (right) in  $D = 5$  dimensions. All quantities are normalized with respect to the mass of the black objects.

However, a different pattern is found for  $D = 6$  solutions. There the charged “fat” BRs exhibit a different limiting behaviour; similar to the vacuum case, they end in a critical merger configuration [4], where a branch of “pinched” BHs is approached in a horizon topology changing transition<sup>10</sup>. The results in [6] together with (27), (28) show that the critical merger EMd

<sup>9</sup>It is interesting to note that, similar to the vacuum case, the reduced angular momentum  $j$  is bounded from above for charged MP BHs with  $k + 1$  equal magnitude angular momenta in  $D = 2k + 5$  only.

<sup>10</sup>The “pinched” BHs possess a spherical horizon topology and can also be studied within the framework in Section 2. Such solutions have been constructed in [6] (in the vacuum case), branching off from a critical MP solution along the stationary zero-mode perturbation of the Gregory-Laflamme-like instability [22, 23].




Figure 6: Same as Figure 5 for  $D = 6$  solutions.

solution has a finite, nonzero area, while the temperature stays also finite and nonzero.

The (area-temperature-charge) diagram of the MP, BRs and black ringoids is shown in Figures 5-7 (in principle, the equation of state  $T(A_H, Q_E)$  can be deduced from there). One can notice that the five dimensional case is special, since, as  $q \rightarrow 1$ ,  $t_H \rightarrow t_H^{(0)} \geq 0$  for  $D = 5$ , while  $t_H \rightarrow 0$  for  $D > 5$ .




Figure 7: Same as Figure 5 for  $D = 7$  charged Myers-Perry black holes (left) and black ringoids (right).

Finally, let us mention that, for any event horizon topology, the gyromagnetic ratio (23) has a remarkable simple expression in terms of  $\alpha$  only<sup>11</sup>,

$$g = D - 3 + \frac{1}{\cosh^2 \alpha}, \quad (32)$$

<sup>11</sup> Note that this is consistent with the general results obtained in [24].

and varies between  $D - 3$  (for maximally charged solutions  $q = 1$ , *i.e.*  $\alpha \rightarrow \infty$ ) and  $D - 2$  (for solutions with an infinitesimally small charge  $q \rightarrow 0$ , *i.e.*  $\alpha \rightarrow 0$ ).

## 4 Further remarks

Fifteen years after the discovery of the BR by Emparan and Reall [2], [3], the study of BHs with a non-spherical horizon topology continues to be a source of excitement in higher dimensional General Relativity. However, most of the black objects with a nonspherical horizon topology studied in the literature describe vacuum configurations only<sup>12</sup>. Moreover, it is worth noticing that even for the case of an event horizon with spherical topology very few solutions with matter fields are known in closed form (for example, the higher dimensional generalization of the Kerr-Newman solution is only known numerically [17], [25], [26], [27]).

The main purpose of this work was to generalize the non-perturbative framework used in [1] for the study of several classes of vacuum black objects with  $k + 1$  equal angular momenta, to the case of Einstein-Maxwell-dilaton theory. Our results show that, similar to the pure Einstein gravity case, for general dilaton coupling constant the problem reduces to solving a set of coupled PDEs with suitable boundary conditions on a rectangular domain, employing an adequate numerical scheme [1].

As a preliminary step before considering the generic case, we have studied solutions of EMD theory with the Kaluza-Klein value of the dilaton coupling constant. In this special limit, the action in  $D$  dimensions is obtained by reducing the  $D + 1$  dimensional vacuum Einstein action, while the solutions are found by embedding the  $D$  dimensional vacuum solutions in  $D + 1$  dimensions and boosting in the extra direction.

The resulting EMD solutions are asymptotically flat, and either possess a regular horizon of spherical topology (and thus represent charged generalizations of MP BHs), or an  $S^{m+1} \times S^{2k+1}$  topology (and thus represent charged BRs and black ringoids). These black objects are characterized by their global charges: their mass, their  $k + 1$  equal magnitude angular momenta, and their electric charge.

As mentioned above, these results were obtained only for a particular value of the dilaton coupling constant<sup>13</sup>. It remains a challenge to generalize such solutions to arbitrary values of the dilaton coupling constant, including the pure Einstein-Maxwell case. The construction of more general configurations (*e.g.* higher dimensional generalizations of the  $D = 5$  dipole BRs [29], solutions with a Chern-Simons term or black objects coupled with a  $p$ -form field (with  $p > 2$ )) is another important open question, just like the inclusion of a cosmological constant. We hope to return elsewhere with a systematic study of these aspects.

---

<sup>12</sup>Higher-dimensional rotating BHs in Einstein gravity coupled to a 2-form or 3-form field strength and to a dilaton with arbitrary coupling have been studied in [19]. These solutions are constructed within the blackfold approach and describe charged MP BHs and various black objects with a non-spherical horizon topology.

<sup>13</sup>Note that an extension of the generating technique in Section 3 can be used to construct (toroidally compactified) heterotic string theory generalizations of the vacuum black objects within the Ansatz (5). In that case, an approach to obtain the charged solutions from the neutral ones was presented in Ref. [28]. Again, the properties of the new configurations can be derived from the corresponding vacuum solutions.

# Acknowledgements

E.R. gratefully acknowledges funding from the FCT-IF programme. This work was partially supported by the H2020-MSCA-RISE-2015 Grant No. StronGrHEP-690904, by the CIDMA project UID/MAT/04106/2013 and by the DFG Research Training Group 1620 “Models of Gravity”.

# References

- [1] B. Kleihaus, J. Kunz and E. Radu, JHEP **1501** (2015) 117 [arXiv:1410.0581 [gr-qc]].
- [2] R. Emparan and H. S. Reall, Phys. Rev. Lett. **88** (2002) 101101 [arXiv:hep-th/0110260].
- [3] R. Emparan and H. S. Reall, Phys. Rev. D **65** (2002) 084025 [arXiv:hep-th/0110258].
- [4] R. Emparan, T. Harmark, V. Niarchos, N. A. Obers and M. J. Rodriguez, JHEP **0710** (2007) 110 [arXiv:0708.2181 [hep-th]].
- [5] B. Kleihaus, J. Kunz and E. Radu, Phys. Lett. B **718** (2013) 1073 [arXiv:1205.5437 [hep-th]].
- [6] Ó. J. C. Dias, J. E. Santos and B. Way, JHEP **1407** (2014) 045 [arXiv:1402.6345 [hep-th]].
- [7] R. C. Myers and M. J. Perry, Annals Phys. **172** (1986) 304.
- [8] R. P. Kerr, Phys. Rev. Lett. **11** (1963) 237.
- [9] B. Kleihaus and J. Kunz, arXiv:1603.07267 [gr-qc].
- [10] H. S. Reall, Int. J. Mod. Phys. D **21** (2012) 1230001 [arXiv:1210.1402 [gr-qc]].
- [11] R. Emparan and H. S. Reall, Living Rev. Rel. **11** (2008) 6 [arXiv:0801.3471 [hep-th]].
- [12] R. Emparan and H. S. Reall, Class. Quant. Grav. **23** (2006) R169 [hep-th/0608012].
- [13] B. Kleihaus, J. Kunz and K. Schnülle, Phys. Lett. B **699** (2011) 192 [arXiv:1012.5044 [hep-th]].
- [14] B. Kleihaus, J. Kunz and E. Radu, Phys. Lett. B **723** (2013) 182 [arXiv:1303.2190 [gr-qc]].
- [15] B. Kleihaus, J. Kunz and E. Radu, Phys. Lett. B **678** (2009) 301 [arXiv:0904.2723 [hep-th]].
- [16] B. Kleihaus, J. Kunz, E. Radu and M. J. Rodriguez, JHEP **1102** (2011) 058 [arXiv:1010.2898 [gr-qc]].
- [17] J. Kunz, F. Navarro-Lerida and A. K. Petersen, Phys. Lett. B **614** (2005) 104 [gr-qc/0503010].
- [18] J. Kunz, D. Maison, F. Navarro-Lerida and J. Viebahn, Phys. Lett. B **639** (2006) 95 [hep-th/0606005].
- [19] M. M. Caldarelli, R. Emparan and B. Van Pol, JHEP **1104** (2011) 013 [arXiv:1012.4517 [hep-th]].

- [20] H. K. Kunduri and J. Lucietti, Phys. Lett. B **609** (2005) 143 [hep-th/0412153].
- [21] S. Grunau, Phys. Rev. D **90** (2014) 064022 [arXiv:1407.2009 [gr-qc]].
- [22] O. J. C. Dias, P. Figueras, R. Monteiro, J. E. Santos and R. Emparan, Phys. Rev. D **80** (2009) 111701 [arXiv:0907.2248 [hep-th]].
- [23] O. J. C. Dias, P. Figueras, R. Monteiro and J. E. Santos, Phys. Rev. D **82** (2010) 104025 [arXiv:1006.1904 [hep-th]].
- [24] M. Ortaggio and V. Pravda, JHEP **0612** (2006) 054 [gr-qc/0609049].
- [25] J. Kunz, F. Navarro-Lerida and J. Viebahn, Phys. Lett. B **639** (2006) 362 [hep-th/0605075].
- [26] J. L. Blazquez-Salcedo, J. Kunz and F. Navarro-Lerida, Phys. Lett. B **727** (2013) 340 [arXiv:1309.2088 [gr-qc]].
- [27] J. L. Blazquez-Salcedo, J. Kunz and F. Navarro-Lerida, Phys. Rev. D **89** (2014) 024038 [arXiv:1311.0062 [gr-qc]].
- [28] A. Sen, Nucl. Phys. B **440** (1995) 421 [arXiv:hep-th/9411187].
- [29] R. Emparan, JHEP **0403** (2004) 064 [hep-th/0402149].